

Reporting Title: Vancomycin, Trough, S
Performing Location: Rochester

Ordering Guidance:

- In addition to this trough assay, both peak level and random testing are available.
1. Serum for a peak level should be drawn 1 hour after completion of dose; order VANPA / Vancomycin, Peak, Serum.
 2. Serum for random testing should be ordered as VANRA / Vancomycin, Random, Serum.

Specimen Requirements:

Collection Container/Tube: Red top
Submission Container/Tube: Plastic vial
Specimen Volume: 0.5 mL

Collection Instructions:

1. Draw specimen immediately prior to the next dose (within 30 minutes).
2. Centrifuge and aliquot serum into a plastic vial within 2 hours of collection.

Forms:

If not ordering electronically, complete, print, and send a [Therapeutics Test Request](#) (T831) with the specimen.

Specimen Type	Temperature	Time	Special Container
Serum Red	Ambient	48 hours	
	Refrigerated (preferred)	14 days	
	Frozen	365 days	

Result Codes:

Result ID	Reporting Name	Type	Unit	LOINC®
VANTA	Vancomycin, Trough, S	Numeric	mcg/mL	4092-3

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

80202

Reference Values:

Therapeutic: 10.0-20.0 mcg/mL