

Reporting Title: Tissue Transglutaminase Ab, IgG, S
Performing Location: Mayo Clinic Laboratories - Rochester Superior Drive

Ordering Guidance:
Cascade testing is recommended for celiac disease. Cascade testing ensures that testing proceeds in an algorithmic fashion. The following cascades are available; select the appropriate one for your specific patient situation.

- CDCOM / Celiac Disease Comprehensive Cascade, Serum and Whole Blood: complete testing including HLA DQ
- CDSP / Celiac Disease Serology Cascade, Serum: complete serology testing excluding HLA DQ
- CDGF / Celiac Disease Gluten-Free Cascade, Serum and Whole Blood: for patients already adhering to a gluten-free diet

To order individual tests, see [Celiac Disease Diagnostic Testing Algorithm](#).

Specimen Requirements:
Supplies: Sarstedt Aliquot Tube 5 mL (T914)
Collection Container/Tube:
Preferred: Serum gel
Acceptable: Red top
Submission Container/Tube: Plastic vial
Specimen Volume: 0.5 mL
Collection Instructions: Centrifuge and aliquot serum into a plastic vial.

Forms:
If not ordering electronically, complete, print, and send a [Gastroenterology and Hepatology Test Request](#) (T728) with the specimen.

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	21 days	
	Frozen	21 days	

Result Codes:

Result ID	Reporting Name	Type	Unit	LOINC®
TTGG	Tissue Transglutaminase Ab, IgG, S	Numeric	U/mL	56537-4

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:
No

CPT Code Information:
86364

Reference Values:
<6.0 U/mL (negative)
6.0-9.0 U/mL (weak positive)

>9.0 U/mL (positive)
Reference values apply to all ages.