

Overview

Useful For

Preferred screening test for detection of arsenic exposure using 24-hour urine specimens

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
SPASU	Arsenic Speciation, 24 Hr, U	Yes	No

Testing Algorithm

If total arsenic concentration is 10 mcg/L or greater, then speciation will be performed at an additional charge.

Special Instructions

- [Urine Preservatives-Collection and Transportation for 24-Hour Urine Specimens](#)
- [Metals Analysis Specimen Collection and Transport](#)

Method Name

Triple-Quadrupole Inductively Coupled Plasma Mass Spectrometry (ICP-MS/MS)

NY State Available

Yes

Specimen

Specimen Type

Urine

Necessary Information

24-Hour volume (in milliliters) is required.

Specimen Required

Patient Preparation:

- For 48-hour period prior to start of collection, patient should not eat seafood.
- High concentrations of gadolinium and iodine are known to interfere with most metal tests. If either gadolinium- or iodine-containing contrast media has been administered, a specimen should not be collected for 96 hours.

Supplies: Urine Tubes, 10 mL (T068)

Collection Container/Tube: Clean, plastic urine container with no metal cap or glued insert.

Submission Container/Tube: Plastic, 10-mL urine tube or clean, plastic aliquot container with no metal cap or glued insert.

Specimen Volume: 6 mL

Collection Instructions:

1. Collect urine for 24 hours.
2. Refrigerate specimen within 4 hours of completion of 24-hour collection.
3. See [Metals Analysis Specimen Collection and Transport](#) for complete instructions.

Additional Information: See [Urine Preservatives-Collection and Transportation for 24-Hour Urine Specimens](#).

Urine Preservative Collection Options

Note: The addition of preservative or application of temperature controls **must occur within 4 hours of completion** of the collection.

Ambient (no additive)	OK
Refrigerate (no additive)	Preferred
Frozen (no additive)	OK
50% Acetic Acid	OK
Boric Acid	No
Diazolidinyl Urea	No
6M Hydrochloric Acid	OK
6M Nitric Acid	OK
Sodium Carbonate	No
Thymol	No
Toluene	No

Specimen Minimum Volume

3 mL

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	28 days	
	Frozen	28 days	
	Ambient	72 hours	

Clinical & Interpretive

Clinical Information

Arsenic (As) is a naturally occurring element that is widely distributed in the Earth's crust. Arsenic is classified chemically as a metalloid, having both metal and nonmetal properties. Elemental arsenic is a steel gray solid material. However,

arsenic is usually found in the environment combined with other elements such as oxygen, chlorine, and sulfur. Arsenic combined with these elements is called inorganic arsenic. Arsenic combined with carbon and hydrogen is referred to as organic arsenic. The organic forms (eg, arsenobetaine and arsenocholine) are relatively nontoxic, while the inorganic forms are toxic. The toxic inorganic forms are arsenite ($\text{As}[3+]/\text{As}[\text{III}]$) and arsenate ($\text{As}[5+]/\text{As}[\text{V}]$). Inorganic As(V) is readily reduced to inorganic As(III), which is then primarily broken down to the less toxic methylated metabolites monomethylarsonic acid and, subsequently, dimethylarsinic acid.

In the past, inorganic arsenic compounds were predominantly used as pesticides, primarily on cotton fields and in orchards. Inorganic arsenic compounds can no longer be used in agriculture. However, organic arsenic compounds, namely cacodylic acid, disodium methylarsenate, and monosodium methylarsenate, are still used as pesticides, principally on cotton. Some organic arsenic compounds are used as additives in animal feed. Small quantities of elemental arsenic are added to other metals to form metal mixtures or alloys with improved properties. The greatest use of arsenic in alloys is in lead-acid batteries for automobiles. Another important use of arsenic compounds is in semiconductors and light-emitting diodes.

People are exposed to arsenic by eating food, drinking water, or breathing air. Of these, food is usually the largest source of arsenic. The predominant dietary source of arsenic is seafood, followed by rice/rice cereal, mushrooms, and poultry. While seafood contains the greatest amounts of arsenic, for fish and shellfish, this is mostly in an organic form of arsenic called arsenobetaine, which is much less harmful. Some seaweed may contain arsenic in the inorganic form, which is more toxic. In the United States, some areas also contain high natural levels of arsenic in rock, which can lead to elevated levels in the soil and drinking water. Occupational (eg, copper or lead smelting, wood treating, or pesticide application) exposure is another source where people may be introduced to elevated levels of arsenic. Lastly, hazardous waste sites may contain large quantities of arsenic and, if not disposed of properly, may get into the surrounding water, air, or soil.

A wide range of signs and symptoms may be seen in acute arsenic poisoning, including headache, nausea, vomiting, diarrhea, abdominal pain, hypotension, fever, hemolysis, seizures, and mental status changes. Symptoms of chronic poisoning, also called arseniasis, are mostly insidious and nonspecific. The gastrointestinal tract, skin, and central nervous system are usually involved. Nausea, epigastric pain, colic abdominal pain, diarrhea, and paresthesias of the hands and feet can also occur.

Since arsenic is excreted predominantly by glomerular filtration, measurement of arsenic in urine is the most reliable means of detecting arsenic exposures within the last several days.

Reference Values

0-17 years: Not established

> or =18 years: <35 mcg/24 h

Interpretation

Physiologically, arsenic exists in toxic and nontoxic forms. The total arsenic concentration reflects all arsenic present in the sample regardless of species (eg, inorganic vs. methylated vs. organic arsenic). The measurement of urinary total arsenic levels is generally accepted as the most reliable indicator of recent arsenic exposure. However, if the total urine arsenic concentration is elevated, arsenic speciation must be performed to identify if it is a toxic form (eg, inorganic and methylated forms) or a relatively nontoxic organic form (eg, arsenobetaine and arsenocholine).

The inorganic toxic forms of arsenic (eg, As(III) and As(V)) are found in the urine shortly after ingestion, whereas the less

toxic methylated forms (monomethylarsonic acid [MMA], dimethylarsinic acid [DMA]) are the species that predominate longer than 24 hours after ingestion. In general, urinary As(III) and As(V) concentrations peak in the urine at approximately 10 hours and return to normal 20 to 30 hours after ingestion. Urinary MMA and DMA concentrations normally peak at approximately 40 to 60 hours and return to baseline 6 to 20 days after ingestion.

After a seafood meal (seafood generally contains the nontoxic, organic form of arsenic (eg, arsenobetaine), the urine output of arsenic may increase to over 300 mcg/24 h, after which it will decline.

This test can determine if the patient has been exposed to above-average levels of arsenic. It cannot predict whether the arsenic levels in their body will affect their health.

Cautions

Consumption of seafood before collection of a urine specimen for arsenic testing is likely to result in a report of an elevated concentration of arsenic found in the urine, which can be clinically misleading.

Clinical Reference

1. Fillol CC, Dor F, Labat L, et al. Urinary arsenic concentrations and speciation in residents living in an area with naturally contaminated soils. *Sci Total Environ*. 2010;408(5):1190-1194
2. Caldwell K, Jones R, Verdon C, et al. Levels of urinary total and speciated arsenic in the US population: National Health and Nutrition Examination Survey 2003-2004. *J Expo Sci Environ Epidemiol*. 2009;19(1):59-68
3. Agency for Toxic Substances and Disease Registry: Toxicological profile for arsenic. US Department of Health and Human Services. Updated August 2007. Accessed August 30, 2024. Available at www.atsdr.cdc.gov/ToxProfiles/tp2.pdf
4. Strathmann FG, Blum LM. Toxic elements. In: Rifai N, Chiu RWK, Young I, Burnham CD, Wittwer CT, eds. *Tietz Textbook of Laboratory Medicine*. 7th ed. Elsevier; 2023:chap 44
5. Keil DE, Berger-Ritchie J, McMillin GA. Testing for toxic elements: A focus on arsenic, cadmium, lead, and mercury. *Lab Med*. 2011;42(12):735-742
6. Navas-Acien A, Francesconi KA, Silbergeld EK, Guallar E. Seafood intake and urine concentrations of total arsenic, dimethylarsinate and arsenobetaine in the US population. *Environ Res*. 2011;111(1):110-8. doi:10.1016/j.envres.2010.10.009
7. Tchounwou PB, Yedjou CG, Udensi UK, et al. State of the science review of the health effects of inorganic arsenic: Perspectives for future research. *Environ Toxicol*. 2019;34(2):188-202. doi:10.1002/tox.22673

Performance

Method Description

The metal analytes of interest are analyzed by triple-quadrupole inductively coupled plasma mass spectrometry.(Unpublished Mayo method).

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days

Specimen Retention Time

14 days

Performing Laboratory Location

Rochester

Fees & Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

82175

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
ASU24	Arsenic w/Reflex, 24 Hr, U	5587-1

Result ID	Test Result Name	Result LOINC® Value
TIME3	Collection Duration	13362-9
VL14	Urine Volume	3167-4
48537	Arsenic, 24 Hr, U	5587-1
92214	Total Arsenic Concentration	21074-0