

Overview

Useful For

Evaluating patients with possible peroxisomal disorders, including peroxisomal biogenesis disorders, X-linked adrenoleukodystrophy, and Refsum disease using plasma specimens

Aiding in the assessment of peroxisomal function

Testing Algorithm

For more information see:

- [Newborn Screen Follow-up for X-Linked Adrenoleukodystrophy](#)
- [Epilepsy: Unexplained Refractory and/or Familial Testing Algorithm](#)

If the patient has abnormal newborn screening results for X-linked adrenoleukodystrophy, refer to the appropriate American College of Medical Genetics and Genomics Newborn Screening ACT Sheet.(1)

Special Instructions

- [Biochemical Genetics Patient Information](#)
- [Newborn Screen Follow-up for X-Linked Adrenoleukodystrophy](#)
- [Epilepsy: Unexplained Refractory and/or Familial Testing Algorithm](#)

Highlights

This test analyzes very long-chain fatty acids as well as pristanic and phytanic acid to aid in diagnosis of peroxisomal biogenesis disorders, X-linked adrenoleukodystrophy (X-ALD), and Refsum disease.

Reports include concentrations of C22:0, C24:0, C26:0 species, phytanic and pristanic acid, and calculated C24:0/C22:0, C26:0/C22:0 and phytanic acid: pristanic acid ratios.

This test is also appropriate for follow-up of an abnormal newborn screen for X-ALD.

Method Name

Gas Chromatography Mass Spectrometry (GC-MS)

NY State Available

Yes

Specimen

Specimen Type

Plasma

Necessary Information

1. Patient's age and sex is required.
2. [Biochemical Genetics Patient Information](#) (T602) is recommended, but not required, to be filled out and sent with the specimen to aid in the interpretation of test results.

Specimen Required

Patient Preparation:

1. Fasting-overnight (12-14 hours). If fasting not possible for babies or infants, collect specimen prior to next feeding.
2. Patient must not consume any alcohol for 24 hours before the specimen is collected.

**Supplies:** Sarstedt Aliquot Tube, 5 mL (T914)

Collection Container/Tube:

**Preferred:** Green top (sodium heparin)

**Acceptable:** Lavender top (EDTA) or green top (lithium heparin)

**Submission Container/Tube:** Plastic vial

**Specimen Volume:** 0.5 mL

**Collection Instructions:** Centrifuge and aliquot plasma into plastic vial.

Forms

1. [Biochemical Genetics Patient Information](#) (T602)
2. If not ordering electronically, complete, print, and send a [Biochemical Genetics Test Request](#) (T798) with the specimen.

Specimen Minimum Volume

0.15 mL

Reject Due To

Gross hemolysis	OK
Gross lipemia	Reject
Gross icterus	OK

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Plasma	Refrigerated	15 days	
	Frozen (preferred)	92 days	

Clinical & Interpretive

Clinical Information

Peroxisomes are organelles present in all human cells except mature erythrocytes. They carry out essential metabolic functions, including beta-oxidation of very long-chain fatty acids (VLCFA), alpha-oxidation of phytanic acid, and

biosynthesis of plasmalogen and bile acids. Peroxisomal disorders include disorders of peroxisomal biogenesis with defective assembly of the entire organelle, and single peroxisomal enzyme/transporter defects where the organelle is intact but a specific function is disrupted. Peroxisomal beta-oxidation of VLCFA is impaired in all disorders of peroxisomal biogenesis and in selected single enzyme deficiencies, particularly X-linked adrenoleukodystrophy (X-ALD), resulting in elevated concentrations of VLCFA in plasma or serum.

Peroxisomal biogenesis disorders (PBD) include Zellweger syndrome spectrum disorders, which are clinically diverse and range in severity from neonatal lethal (Zellweger syndrome) to more variable clinical courses in neonatal adrenoleukodystrophy and infantile Refsum disease. Affected children typically have hypotonia, poor feeding, distinctive facial features, seizures, and liver dysfunction. Other features can include retinal dystrophy, hearing loss, developmental delays, and bleeding episodes. Rhizomelic chondrodysplasia punctata is another PBD. It is characterized by rhizomelic shortening, chondrodysplasia punctata, cataracts, intellectual disability, and seizures, although it can have a milder phenotype with only cataracts and chondrodysplasia. The typical biochemical profile shows normal VLCFA and elevated phytanic acid.

X-ALD is a neurologic disorder affecting the white matter and adrenal cortex. It can present between ages 4 and 8 years as a childhood cerebral form with behavioral and cognitive changes, associated with neurologic decline. Other forms include an "Addison disease only" phenotype with adrenocortical insufficiency without initial neurologic abnormality and adrenomyeloneuropathy associated with later-onset progressive paraparesis. X-ALD is an X-linked condition that primarily affects male patients; however, some female patients who are carriers can develop later-onset neurologic manifestations. In 2016, X-ALD was added to the US Recommended Uniform Screening Panel, a list of conditions that are nationally recommended for newborn screening by the Secretary's Advisory Committee on Heritable Disorders in Newborns and Children.

Refsum disease is a peroxisomal disorder characterized by anosmia, retinitis pigmentosa, neuropathy, deafness, ataxia, ichthyosis, and cardiac abnormalities. The classic biochemical profile of Refsum disease is an elevated plasma or serum phytanic acid level.

Biochemical abnormalities in peroxisomal disorders include accumulations of VLCFA, phytanic acid, and pristanic acid. The differential diagnosis of these disorders is based on recognition of clinical phenotypes combined with a series of biochemical tests to assess peroxisomal function and structure. These include measurements and ratios of VLCFA, pipelicolic acid (PIPA / Pipecolic Acid, Serum; PIPU / Pipecolic Acid, Random, Urine), phytanic acid and its metabolite pristanic acid. In addition, confirmatory testing for X-ALD (XALDZ / X-Linked Adrenoleukodystrophy, Full Gene Analysis, Varies) via molecular genetic analysis is available.

### Reference Values

C22:0

< or =96.3 nmol/mL

C24:0

< or =91.4 nmol/mL

C26:0

< or =1.30 nmol/mL

**C24:0/C22:0 RATIO**

&lt; or =1.39

**C26:0/C22:0 RATIO**

&lt; or =0.023

**PRISTANIC ACID**

0-4 months: &lt; or =0.60 nmol/mL

5-8 months: &lt; or =0.84 nmol/mL

9-12 months: &lt; or =0.77 nmol/mL

13-23 months: &lt; or =1.47 nmol/mL

&gt; or =24 months: &lt; or =2.98 nmol/mL

**PHYTANIC ACID**

0-4 months: &lt; or =5.28 nmol/mL

5-8 months: &lt; or =5.70 nmol/mL

9-12 months: &lt; or =4.40 nmol/mL

13-23 months: &lt; or =8.62 nmol/mL

&gt; or =24 months: &lt; or =9.88 nmol/mL

**PRISTANIC/PHYTANIC ACID RATIO**

0-4 months: &lt; or =0.35

5-8 months: &lt; or =0.28

9-12 months: &lt; or =0.23

13-23 months: &lt; or =0.24

&gt; or =24 months: &lt; or =0.39

**Interpretation**

Reports include concentrations of C22:0, C24:0, C26:0 species, phytanic acid and pristanic acid, and calculated C24:0/C22:0, C26:0/C22:0 and phytanic acid:pristanic acid ratios. When no significant abnormalities are detected, a simple descriptive interpretation is provided.

A profile of elevated phytanic acid, low-normal pristanic acid, and normal very long-chain fatty acids is suggestive of Refsum disease (phytanic acid oxidase deficiency); however, phytanic acid concentration may also be increased in disorders of peroxisomal biogenesis and should be considered in the differential diagnosis of peroxisomal disorders.

If results are suggestive of hemizygoty for X-linked adrenoleukodystrophy, the calculated value of a discriminating function that more accurately segregates hemizygous individuals from normal controls is included in the report.

Positive test results could be due to a genetic or nongenetic condition. Additional confirmatory testing would be required to differentiate between these causes.

**Cautions**

In rare instances, patients with X-linked adrenoleukodystrophy (X-ALD) may have only minimally elevated values; 15% to 20% of women heterozygous for X-ALD have normal plasma very long-chain fatty acid levels.

False-positive results may occur with nonfasting specimens.

### Clinical Reference

1. Newborn Screening ACT Sheet [Elevated Lysophosphatidylcholine] X-Linked Adrenoleukodystrophy (X-ALD). American College of Medical Genetics and Genomics; 2023. Revised November 2023. Accessed March 25, 2025. Available at [www.acmg.net/PDFLibrary/X-ALD-ACT-Sheet.pdf](http://www.acmg.net/PDFLibrary/X-ALD-ACT-Sheet.pdf)
2. Moser AB, Kreiter N, Bezman L, et al. Plasma very long chain fatty acid assay in 3,000 peroxisome disease patients and 29,000 controls. *Ann Neurol*. 1999;45:100-110
3. Turk BR, Theda C, Fatemi A, Moser AB. X-linked adrenoleukodystrophy: Pathology, pathophysiology, diagnostic testing, newborn screening and therapies. *Int J Dev Neurosci*. 2020;80(1):52-72. doi:10.1002/jdn.10003
4. Waterham HR, Ferdinandusse S, Wanders RJA. Human disorders of peroxisome metabolism and biogenesis. *Biochimica et Biophysica Acta*. 2016;1863(5):922-933. doi:10.1016/j.bbamcr.2015.11.015

### Performance

#### Method Description

Acidic hydrolysis is followed by basic hydrolysis and reacidification. Hexane extraction then proceeds to derivatization with pentafluorobenzyl bromide (PFB). Separation and detection of PFB-esters is accomplished by capillary gas chromatography mass spectrometry using electron capture ionization and selected negative ion monitoring. Quantitation is enhanced by the use of stable isotope-labeled internal standards.(Stellard F, ten Brink HJ, Kok RM, et al. Stable isotope dilution analysis of very long chain fatty acids in plasma, urine and amniotic fluid by electron capture negative ion mass fragmentography. *Clin Chim Acta*. 1990;192:133-144; Rattay TW, Rautenberg M, Sohn AS, et al. Defining diagnostic cutoffs in neurological patients for serum very long chain fatty acids (VLCFA) in genetically confirmed X-adrenoleukodystrophy. *Sci Rep*. 2020;10[1]:15093)

#### PDF Report

No

#### Day(s) Performed

Monday through Friday

#### Report Available

3 to 5 days

#### Specimen Retention Time

1 month

#### Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

82726

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
POXP	Fatty Acid Profile, Peroxisomal,P	43677-4

Result ID	Test Result Name	Result LOINC® Value
36641	C22:0	30194-5
36642	C24:0	30195-2
36643	C26:0	30197-8
36644	C24:0/C22:0	30196-0
36645	C26:0/C22:0	30198-6
36646	Pristanic Acid	22761-1
36647	Phytanic Acid	22671-2
36648	Pristanic/Phytanic	30550-8
36649	Interpretation (POXP)	59462-2