

Risankizumab, Serum

Test ID: RISA

Useful for:

Assessing the response to risankizumab therapy

Assessing the need for dose escalation

Evaluating potential changes or discontinuation of therapy

Monitoring patients who need to be above a certain risankizumab concentration to improve the odds of a clinical response for therapy optimization

Methods:

Liquid Chromatography Mass Spectrometry (LC-MS)

Reference Values:

Lower limit of quantitation = 1.0 mcg/mL

Specimen Requirements:

Supplies: Sarstedt Aliquot Tube 5 mL (T914)

Container/Tube:

Preferred: Red top

Acceptable: Serum gel

Submission Container/Tube: Plastic vial

Specimen Volume: 0.5mL

Collection Instructions:

1. Draw blood immediately before the next scheduled dose (trough specimen).
2. Centrifuge and aliquot serum into plastic vial within 2 hours of collection.

Minimum Volume: 0.25mL

Specimen Stability Information:

| Specimen Type | Temperature | Time |
|---------------|--------------------------|---------|
| Serum | Refrigerated (preferred) | 28 days |
| | Ambient | 28 days |
| | Frozen | 28 days |

Cautions:

Lipemic samples will be rejected.

CPT Code:

80299

Day(s) Performed: Wednesday

Report Available: 2 to 9 days

Questions

Contact Amy Ennis, Laboratory Resource Coordinator at 800-533-1710.